

Four Generation CP Violation in $B \rightarrow \phi K^0, \pi^0 K^0, \eta' K^0$ and Hadronic Uncertainties

Wei-Shu Hou^{a,*}, Makiko Nagashima^{a,†}, Guy Raz^{b,‡} and Andrea Soddu^{b,§}

^a*Department of Physics, National Taiwan University, Taipei, Taiwan 10617, R.O.C.*

^b*Department of Particle Physics, Weizmann Institute of Science, Rehovot 76100, Israel*

(Dated: February 2, 2008)

The fourth generation can give the correct trend of $\mathcal{S}_{\phi K^0}, \mathcal{S}_{\pi^0 K^0} < \sin 2\phi_1/\beta$, as indicated by data, and the effect, being largely leading order, is robust against hadronic uncertainties. The effect on $\mathcal{S}_{\eta' K^0}$, however, is diluted away by hadronic effects, and $\mathcal{S}_{\eta' K^0} \simeq \sin 2\phi_1/\beta$ is expected. The near maximal $\arg V_{t's}^* V_{t'b} \lesssim 90^\circ$ that is needed could resolve the unequal direct CP violation seen in $B \rightarrow K^+ \pi^-$ and $K^+ \pi^0$ modes, and is consistent with $b \rightarrow s \ell^+ \ell^-$ and B_s mixing constraints.

PACS numbers: 11.30.Er, 11.30.Hv, 12.60.Jv, 13.25.Hw

At present there are two hints for possible New Physics (NP) from CP violation (CPV) studies in the B system, both in charmless $b \rightarrow s$ transitions.

Time dependent CPV (TCPV) in B decays to CP eigenstate f is measured by

$$\frac{\Gamma(\bar{B}^0(t) \rightarrow f) - \Gamma(B^0(t) \rightarrow f)}{\Gamma(\bar{B}^0(t) \rightarrow f) + \Gamma(B^0(t) \rightarrow f)} = S_f \sin(\Delta m_B t) - C_f \cos(\Delta m_B t). \quad (1)$$

We concern ourselves with S_f only, since $-C_f = A_f$ are all consistent with zero so far. The current world average of S_f in $b \rightarrow c\bar{c}s$ decays gives $\sin 2\phi_1/\beta = 0.69 \pm 0.03$ [1], which dominantly comes from $B^0 \rightarrow J/\psi K_S$. TCPV measurements in loop dominated $b \rightarrow s\bar{q}q$ processes such as $B^0 \rightarrow \eta' K^0, \phi K^0$ and $\pi^0 K^0$, on the other hand, have persistently given values below $\sin 2\phi_1/\beta$. The current values [1] are $\mathcal{S}_{\eta' K^0} = 0.50 \pm 0.09$ [2], $\mathcal{S}_{\phi K^0} = 0.47 \pm 0.19$ and $\mathcal{S}_{\pi^0 K^0} = 0.31 \pm 0.26$.

It was suggested some time ago [3] that, for $b \rightarrow s\bar{q}q$ final states, a significant $\Delta\mathcal{S}_f = \mathcal{S}_f - \sin 2\phi_1/\beta$ would indicate NP. The study of theoretical uncertainties for $\Delta\mathcal{S}_f$ has therefore been a great focus during the past year. Consensus has emerged that $\Delta\mathcal{S}_f$ in these modes tend to be small and *positive* [4, 5, 6] within the Standard Model (SM), which is opposite the trend seen by experiment. It is therefore imperative to establish $\Delta\mathcal{S}_f \neq 0$ experimentally beyond any doubt in a few modes, which would require considerably more data than present.

A simpler measurement than $\mathcal{S}_f$ is direct CPV (DCPV) asymmetries in flavor-specific final states, which does not require time dependent measurement. DCPV was recently observed [7] in $B^0 \rightarrow K^+ \pi^-$ decay, i.e. $\mathcal{A}_{K^+ \pi^-} = -0.115 \pm 0.018$. Having similar dominating penguin and tree contributions, one would naively expect that $\mathcal{A}_{K^+ \pi^-} = \mathcal{A}_{K^+ \pi^0}$. However, no indication of DCPV was seen in charged $B^+ \rightarrow K^+ \pi^0$, i.e.

$\mathcal{A}_{K^+ \pi^0} = 0.04 \pm 0.04$. The difference with $\mathcal{A}_{K^+ \pi^-}$ could be due to an enhanced color-suppressed amplitude C [8], or electroweak penguin P_{EW} effects [9, 10, 11, 12]. The former requires C to effectively cancel the SM phase in color-allowed tree amplitude T , without recourse to NP. For the latter, NP CPV phases would be needed in P_{EW} .

It would be intriguing if the two hints of NP, one in $\Delta\mathcal{S}_f \neq 0$, the other in $\mathcal{A}_{K^+ \pi^0} - \mathcal{A}_{K^+ \pi^-} \neq 0$, could be manifestations of the same NP source. Since π^0 and ϕ (but not π^-) can materialize from a virtual Z , the $B \rightarrow \pi^0 K^0, \phi K^0$ modes are sensitive to Z penguins. The effect of a NP phase in P_{EW} on $\Delta\mathcal{S}_f$, among several NP scenarios, was studied in Ref. [6]. In another work, some of us have shown [12] that the 4th generation could provide a solution to the $\mathcal{A}_{K^+ \pi^0} - \mathcal{A}_{K^+ \pi^-} \neq 0$ problem through the electroweak penguin. The 4th generation can make specific impact on P_{EW} because the t' quark, like the SM top, enjoys nondecoupling in P_{EW} [13], but largely decouples from photonic and gluonic penguins. Furthermore, it can provide a new CPV phase [14] through $V_{t's}^* V_{t'b} \equiv r_{sb} e^{i\phi_{sb}}$.

In this work we show that the fourth generation can, for the right choice of ϕ_{sb} , give the correct trend for $\Delta\mathcal{S}_f$ in $B^0 \rightarrow \pi^0 K^0$ and ϕK^0 , and is robust against hadronic uncertainties. In contrast, we find $\Delta\mathcal{S}_{\eta' K^0}$ is largely diluted by hadronic effects that are needed to account for the large rate.

For relevant 4th generation parameters, we take [12]

$$m_{t'} = 300 \text{ GeV}, \quad r_{sb} \simeq 0.025, \quad (2)$$

and vary ϕ_{sb} phase. Eq. (2) is consistent with $b \rightarrow s\ell^+ \ell^-$ and B_s mixing constraints [12]. Larger $m_{t'}$ or r_{sb} [15] could lead to larger effects on $\Delta\mathcal{S}_f$, but could run into trouble with the other $b \rightarrow s$ constraints. To study (factorization) model dependence, we compare results in naive factorization (NF) [16], QCD factorization (QCDF) [17, 18] and PQCD [19]. We further use QCDF to illustrate hadronic uncertainties. We choose to use QCDF and PQCD circa 2003 because, in part stimulated by the $\Delta\mathcal{S}$ and $\mathcal{A}_{K^+ \pi^0}$ problems, these factorization models are still being refined.

We adopt QCDF as our reference framework. Defining $\lambda_i \equiv V_{is}^* V_{ib}$, one has $\lambda_u + \lambda_c + \lambda_t + \lambda_{t'} = 0$ with existence of t' . To good approximation, λ_u is negligible compared

*Electronic address: wshou@phys.ntu.edu.tw

†Electronic address: makiko@phys.ntu.edu.tw

‡Electronic address: guy.raz@weizmann.ac.il

§Electronic address: andrea.soddu@weizmann.ac.il

with $\lambda_c \simeq 0.04$, where we have taken the convention to keep V_{cb} real, and placing the 3 CPV phases in V_{ub} , $V_{t's}$ and $V_{t'd}$ [20, 21]. This makes clear correspondence to the standard phase convention for 3 generation case. The unitarity condition $\lambda_t \simeq -\lambda_c - \lambda_{t'}$ allows one to absorb the t effect into the λ_c dependent part (SM term), and the NP $\lambda_{t'}$ dependent part that respects GIM [12, 14].

The $\overline{B}^0 \rightarrow \pi^0 \overline{K}^0$ amplitude is

$$\mathcal{M}_{\pi^0 \overline{K}^0} \propto f_\pi F_{BK} (\lambda_u a_2^u + \frac{3}{2} \lambda_c \alpha_{3,EW}^p - \frac{3}{2} \lambda_{t'} \Delta \alpha_{3,EW}^p) - f_K F_{B\pi} [\lambda_c (\alpha_4^p - \frac{1}{2} \alpha_{4,EW}^p + \beta_3^p) + \frac{1}{2} \lambda_{t'} \Delta \alpha_{4,EW}^p], \quad (3)$$

where $\alpha_{i,(EW)}^p$ and β_3^p are defined in Ref. [17] and evaluated for the $\pi^0 \overline{K}^0$ final state, and $\Delta \alpha_{i,EW}^p$ is the effective (t subtracted) t' contribution. For $\overline{B}^0 \rightarrow \phi \overline{K}^0$, there is no tree term, and one has

$$\mathcal{M}_{\phi \overline{K}^0} \propto \lambda_c [\alpha_3^p + \alpha_4^p + \beta_3^p - \frac{1}{2} (\alpha_{3,EW}^p + \alpha_{4,EW}^p + \beta_{3,EW}^p)] + \frac{\lambda_{t'}}{2} (\Delta \alpha_{3,EW}^p + \Delta \alpha_{4,EW}^p + \Delta \beta_{3,EW}^p), \quad (4)$$

where $\alpha_{i,(EW)}^p$ and β_3^p are evaluated for the $\phi \overline{K}^0$ final state. We have dropped the common $f_\phi F_{BK}$ factor compared to Eq. (4), and we show only the more important terms. The numerics was done with full details according to Ref. [18]. The formula for $\overline{B}^0 \rightarrow \eta' \overline{K}^0$ can be analogously written, but is more elaborate which we do not reproduce here. We stress that the same expressions apply to the amplitudes in NF framework as well, with the various coefficients taken at LO instead of NLO.

In this work we estimate and quantify the impact of hadronic uncertainties in QCDF. Among the hadronic parameters that enter the decay amplitudes, three stand out as having the largest impact due to uncertainties [17]: the divergent part of the hard spectator scattering integral X_H , the divergent part of the weak annihilation integral X_A , and the first inverse moment of the B meson distribution amplitude λ_B . The first two are estimated to be complex numbers of order $\ln(m_B/\Lambda_h)$ with $\Lambda_h = 500$ MeV, and can therefore be parameterized by [22]

$$X_{H,A} = (1 + \rho_{H,A} e^{i\phi_{H,A}}) \ln \frac{m_b}{\Lambda_h}. \quad (5)$$

Our estimate of the hadronic uncertainties is based on the variation of these parameters over a wide range as indicated in Ref. [17]. For reference, we also take as baseline a “standard” scenario, in which we fix $\rho_H = 0$, $\rho_A = 1$, $\phi_A = -45^\circ$ and $\lambda_B = 350$ MeV. This scenario corresponds to the “S3” scenario of Ref. [18], although small numerical differences in input parameters may lead to a slight difference in final results [23].

For $\overline{B}^0 \rightarrow \pi^0 \overline{K}^0$ in PQCD factorization, we adopt the LO result used in Ref. [12]

$$\mathcal{M}_{\pi^0 \overline{K}^0} \propto \lambda_u f_\pi F_{ek}^P + \lambda_c (-f_K F_e^P - f_B F_a^P + f_\pi F_{ek}^P) - \lambda_{t'} f_\pi \Delta F_{ek}^P, \quad (6)$$




FIG. 1: (a) $S_{\pi^0 K^0}$, (b) $S_{\phi K^0}$ vs ϕ_{sb} for QCDF at NLO with “S3” parameters (solid), PQCD at LO (dashed), and NF (dots), which is QCDF at LO. The horizontal solid band is the current experimental range.

where F_e^P , F_a^P , F_{ek} and F_{ek}^P are the strong penguin, strong penguin annihilation, color suppressed tree and (color allowed) electroweak penguin contributions, respectively. These factorizable contributions can be computed by following Ref. [19], and are tabulated in Ref. [12].

For $\overline{B}^0 \rightarrow \phi \overline{K}^0$, we have

$$\mathcal{M}_{\phi \overline{K}^0} \propto \lambda_c (f_\phi F_e^P + f_B F_a^P) - \lambda_{t'} f_\phi \Delta F_e^P, \quad (7)$$

where the F_i^P s are evaluated for $\phi \overline{K}^0$ [24] and not the same as in Eq. (6). We have performed only an approximate computation in this case. We assume that the scale t , where the Wilson coefficients are evaluated, has a mild dependence on the momentum fraction x and the impact parameter b which is conjugate to the parton transverse momentum. The amplitude F_e^P , which is obtained by integrating over the variables x and b , becomes then proportional to $a_e(t)$ with

$$a_e(t) = C_3 + \frac{C_4}{3} + C_4 + \frac{C_3}{3} + C_5 + \frac{C_6}{3} - \frac{1}{2} \left(C_7 + \frac{C_8}{3} + C_9 + \frac{C_{10}}{3} + C_{10} + \frac{C_9}{3} \right). \quad (8)$$

By knowing now the numerical value of F_e^P [19] and the Wilson coefficients in the SM calculated at $t = m_b$, one can then calculate ΔF_e^P with

$$\Delta F_e^P = F_e^P \left(\frac{a_e^{NP}}{a_e^{SM}} - 1 \right). \quad (9)$$

The same procedure is not possible for F_a^P and we keep only the SM contribution by assuming $\Delta F_a^P = 0$ [25]. For $\eta' K$ mode, not much work has been done in PQCD.

To study the model dependence in different factorization approaches, we plot $S_{\pi^0 K^0}$ and $S_{\phi K^0}$ vs ϕ_{sb} in Fig. 1 for QCDF at NLO (with “S3” parameters), PQCD at LO, and NF. The latter is far from realistic (for rates) and is just for comparison. We see that in all three models, $S_{\pi^0 K^0}$ and $S_{\phi K^0}$ dip below $\sin 2\phi_1/\beta$ for $\sin \phi_{sb} \gtrsim 0$, especially around $\phi_{sb} \sim 90^\circ$. Indeed, for a given size of




FIG. 2: (a) $S_{\pi^0 K^0}$, (b) $\mathcal{B}(\pi^0 K^0)$, (c) $S_{\phi K^0}$, (d) $\mathcal{B}(\phi K^0)$, (e) $S_{\eta' K^0}$, (f) $\mathcal{B}(\eta' K^0)$ vs ϕ_{sb} , in QCDF at NLO. The curves are for the “S3” hadronic parameter settings. The light gray regions correspond to varying the hadronic parameters X_A , X_H and λ_B over the range indicated in Eq. (10). The dark gray regions are obtained by varying the hadronic parameters over the same range as above, but keeping only the values for which the branching ratios are within 3σ of the experimental central values.

NP contribution, a choice of a maximal weak phase of 90° (or 270°) tends to maximize the NP effect on CPV while minimizing the NP effect on BR. It is interesting to note that this is precisely what is needed for the 4th generation to help resolve [12] the $\mathcal{A}_{K^+\pi^0} - \mathcal{A}_{K^+\pi^-} \neq 0$ problem. Independently, $\phi_{sb} \sim 90^\circ$ is also the parameter space where $b \rightarrow s\ell^+\ell^-$ and Δm_{B_s} constraints are best evaded [12, 14]. For $\phi_{sb} \sim 270^\circ$, although the $b \rightarrow s\ell^+\ell^-$ and Δm_{B_s} constraints can still be tamed, both $S_{\pi^0 K^0, \phi K^0}$ and $\mathcal{A}_{K^+\pi^0}$ would be in disagreement with experiment.

Hadronic parameters such as strong phases easily affect branching ratios and DCPV asymmetries. Strong phases are definitely present in $B \rightarrow K\pi$ decay as evidenced by the sizable $\mathcal{A}_{K^+\pi^-} = -0.115 \pm 0.018$. The S_f parameter, however, measures the weak phase of the decay amplitude, and is less affected by hadronic parameters [6]. As mentioned, we illustrate this point, by varying the hadronic parameters of QCDF at NLO around the “S3” scenario settings. In particular, we vary [17, 18]

$$\begin{aligned} \rho_{A,H} &\in (0, 1), \quad \phi_{A,H} \in [0, 2\pi], \\ \lambda_B &\in [200, 500] \text{ MeV}, \end{aligned} \quad (10)$$

for the X_A , X_H and λ_B parameters.

We plot $S_{\pi^0 K^0}$, $S_{\phi K^0}$ and $S_{\eta' K^0}$ vs ϕ_{sb} in the left side of Fig. 2. The light shaded regions correspond to varying the parameters over the whole range indicated in




FIG. 3: Ratio of branching ratios (a) R_c and (b) R_n for B^+ and B^0 decay to $K\pi$, as defined in Ref. [9], vs ϕ_{sb} , for QCDF at NLO. The curves and the light gray regions are obtained as in Fig. 2.

Eq. (10). The dark shaded regions correspond to varying the hadronic parameters over the same range, but keeping, for each mode, only the values that produce a branching ratio (right side of Fig. 2) within 3σ of the experimental central value. One sees that, indeed, the branching ratios are strongly affected by the hadronic parameters, and most of the hadronic parameter space cannot survive the bulk of rate and DCPV data when considered together. In contrast, the range of variation for S_f is much more subdued. This is encouraging: the NP effect in S_f for the $\pi^0 K^0$, ϕK^0 and $\eta' K^0$ modes is robust.

We note that the effect of hadronic parameters, when varied over the whole range, is rather strong for $S_{\phi K^0}$. However, when the experimental constraints on the $B \rightarrow \phi K^0$ branching ratio are taken into account, the hadronic uncertainty in $S_{\phi K^0}$ is highly diminished.

Note, also, that $S_{\eta' K^0}$ gets strongly diluted away. The reason behind this is the rather large rate of $B \rightarrow \eta' K$ decay, which seemingly draws from CP conserving (“hadronic”) effects, since there is little evidence for CPV i.e. $\mathcal{A}_{\eta' K} \simeq 0$ [1]. Furthermore, the Z penguin contribution has relatively small strength. We believe the dilution of $S_{\eta' K^0}$ is a generic effect, that is, it is very hard for NP CPV effects to shine through the large hadronic effects, and $S_{\eta' K^0} \simeq \sin 2\phi_1/\beta$ should be expected. In this respect, the Belle result of $S_{\eta' K^0} = 0.62 \pm 0.12 \pm 0.04$, which is fully consistent with $\sin 2\phi_1/\beta = 0.69 \pm 0.03$, is easier to explain in most NP models [2]. If the BaBar result holds out eventually, it would need some conspiracy between NP and hadronic effects to realize theoretically.

We offer some remarks before closing. We have studied the ratio of branching ratios R_c , R_n and R , which are for B^+ , B^0 , and the lifetime corrected $K^+\pi^-$ over $K^0\pi^+$ ratio, respectively. Indeed, these rate ratios are attractive in that they suffer considerably less hadronic uncertainties. We plot R_c and R_n vs ϕ_{sb} in Fig. 3 for QCDF at NLO and varying hadronic parameters over the full range of Eq. (10). The contrast with the branching ratio plot in Fig. 2 is striking. Interestingly, for $|\phi_{sb}| \lesssim 80^\circ$, the hadronic uncertainties are even less significant (a bit more for R), and the results with 4th generation are basically consistent with experiment. But for $\phi_{sb} \sim \pi$, besides much larger hadronic uncertainties, R_c and R_n would deviate substantially (being larger) from

data, and disallowed. This is again consistent with the analysis from $\mathcal{A}_{K^+\pi^0} - \mathcal{A}_{K^+\pi^-}$ as well as $b \rightarrow s\ell^+\ell^-$ and B_s mixing, and with our findings for $S_{\pi K}$ and $S_{\phi K}$.

S_f has been studied experimentally in quite a few other modes such as $f = f_0(980)K_S, \omega K^0$ [1], as well as 3-body modes such as $f = K\bar{K}K$ and $K_S\pi^0\pi^0$. The interest in $S_{\eta K}, S_{\omega K}$ and $S_{\rho^0 K}$ have been stressed [4, 6]. We have studied these modes and found the effect of hadronic uncertainties to be more significant. Thus, experimental studies in these modes would shed little light on NP parameters, except that $S_{\rho^0 K} > \sin 2\phi_1/\beta$ is likely realized. The theory for 3-body modes is even less developed. Similarly, DCPV depends sensitively on hadronic phases, and much theoretical work is currently ongoing to elucidate these. We therefore leave this for future studies. Our studies do show that DCPV in the above mentioned 2-body modes are in general consistent with data, since experimental errors are still large. The only firmly measured DCPV is in $\mathcal{A}_{K^+\pi^-}$, while Ref. [12] has demonstrated that the 4th generation may help resolve the $\mathcal{A}_{K^+\pi^0} - \mathcal{A}_{K^+\pi^-} \neq 0$ problem.

Finally, we note from Fig. 2 that for QCDF the experimental central values are unattainable once the branching ratio is constrained to within 3σ of experiment. (Note, however, from Fig. 1, that our approximate PQCD

result could fit the $S_{\phi K}$ central value.) If the experimental central values for $S_{\phi K}, S_{\pi K}$ and $S_{\eta' K}$ persist, more work on factorization models seem needed to shed further light on whether the 4th generation, or other New Physics, could account for the observed effect.

In summary, we have studied in this work the effect of a 4th generation model on the TCPV parameter S_f for $f = \pi K^0, \phi K^0$ and $\eta' K^0$. We have shown, using QCDF at NLO, that the NP effects on these S_f 's are rather robust against hadronic uncertainties. This robustness may be generic to a large class of NP models. We found that the same 4th generation parameters that explain $\mathcal{A}_{K^+\pi^0} \sim 0$ while $\mathcal{A}_{K^+\pi^-} \simeq -11\%$, can give the correct trend in S_f . However, we also showed that $S_{\eta' K^0}, S_{\pi K^0}$ and to a lesser degree $S_{\phi K^0}$ are predicted to be closer to $\sin 2\phi_1/\beta$ than the current data indicate. Due to the robustness of the S_f , better measurements could provide an important test of the 4th generation model as well as other NP models.

Acknowledgement. This work is supported in part by NSC-94-2112-M-002-035 and NSC94-2811-M-002-053 of Taiwan, and HPRN-CT-2002-00292 of Israel. We would like to thank M. Beneke and Y. Nir for very useful discussions.

-
- [1] Heavy Flavor Averaging Group [HFAG], see webpage <http://www.slac.stanford.edu/xorg/hfag/>.
 - [2] The BaBar value of $S_{\eta' K^0} = 0.36 \pm 0.13 \pm 0.03$, and the Belle value of $S_{\eta' K^0} = 0.62 \pm 0.12 \pm 0.04$ are in some variance. One could inflate the error of the mean by a factor of $\sqrt{2}$. We refrain from doing so and use the value quoted by HFAG, for sake of consistency with other modes. Note that the Belle value is in good agreement with $\sin 2\phi_1$.
 - [3] Y. Grossman and M.P. Worah, Phys. Lett. B **395**, 241 (1997); R. Fleischer, Int. J. Mod. Phys. A **12**, 2459 (1997).
 - [4] M. Beneke, Phys. Lett. B **620**, 143 (2005).
 - [5] H. Y. Cheng, C. K. Chua and A. Soni, Phys. Rev. D **72**, 014006 (2005).
 - [6] G. Buchalla, G. Hiller, Y. Nir and G. Raz, JHEP **0509**, 074 (2005).
 - [7] B. Aubert *et al.* [BaBar Collab.], Phys. Rev. Lett. **93**, 131801 (2004); Y. Chao *et al.* [Belle Collab.], *ibid.* **93**, 191802 (2004); K. Abe *et al.* [Belle Collab.], hep-ex/0507045.
 - [8] C.W. Chiang, M. Gronau, J.L. Rosner and D.A. Suprun, Phys. Rev. D **70**, 034020 (2004); Y.Y. Charng and H.n. Li, *ibid.* D **71**, 014036 (2005); M. Gronau and J.L. Rosner, *ibid.* D **71**, 074019 (2005); C.S. Kim, S. Oh and C. Yu, *ibid.* D **95**, 141601 (2005); H.n. Li, S. Mishima and A.I. Sanda, Phys. Rev. D **72**, 114005 (2005).
 - [9] A.J. Buras, R. Fleischer, S. Recksiegel and F. Schwab, Eur. Phys. J. C **32**, 45 (2003); Phys. Rev. Lett. **92**, 101804 (2004); Nucl. Phys. B **697**, 133 (2004); Acta Phys. Polon. B **36**, 2015 (2005).
 - [10] V. Barger, C.W. Chiang, P. Langacker and H.S. Lee, Phys. Lett. B **598**, 218 (2004).
 - [11] S. Baek *et al.*, Phys. Rev. D **71**, 057502 (2005).
 - [12] W.S. Hou, M. Nagashima and A. Soddu, Phys. Rev. Lett. **95**, 141601 (2005).
 - [13] W.S. Hou, R.S. Willey and A. Soni, Phys. Rev. Lett. **58**, 1608 (1987).
 - [14] A. Arhrib and W.S. Hou, Eur. Phys. J. C **27**, 555 (2003).
 - [15] Choosing a larger $m_{t'}$ can be compensated by a smaller r_{sb} , and vice versa, for the same effect.
 - [16] A. Ali, G. Kramer and C. D. Lu, Phys. Rev. D **58**, 094009 (1998).
 - [17] M. Beneke, G. Buchalla, M. Neubert and C. Sachrajda, Nucl. Phys. B **606**, 245 (2001).
 - [18] M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003).
 - [19] Y.Y. Keum, H.n. Li and A.I. Sanda, Phys. Rev. D **63**, 054008 (2001).
 - [20] W.S. Hou, A. Soni and H. Steger, Phys. Lett. B **192**, 441 (1987).
 - [21] W.S. Hou, M. Nagashima and A. Soddu, Phys. Rev. D **72**, 115007 (2005).
 - [22] In principle both X_H and X_A are mode dependent. However, since we vary them over the whole possible range in any case, this point does not affect our analysis.
 - [23] M. Beneke, private communications.
 - [24] S. Mishima, Phys. Lett. B **521**, 252 (2001); S. Mishima and A.I. Sanda, Prog. Theor. Phys. **110**, 549 (2003).
 - [25] Since one expects very strong GIM cancellation between t' and t for strong penguin, the approximation should be a very good one.